

BABY CALF

ROWE 18% CALF STARTER

R30 SWEET

Starter Feed for Calves up to 200 pounds



Bovamine® Defend™ is a unique living probiotic that stabilizes rumen and small intestine digestive functions while stimulating natural and healthy digestion by reinforcing and enhancing the animal's natural defenses against harmful bacteria.

MEDICATED

For the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT

Monensin 30 g/T

CONTAINS DIFLUBENZURON

An insect growth regulator, which prevents the development of house, stable, face, and horn flies in the manure of treated cattle.

Diflubenzuron..... 3.65 Gms/Ton

GUARANTEED ANALYSIS

Crude Protein, Minimum.....	18.00%
Crude Fat, Minimum.....	2.50%
Crude Fiber, Maximum.....	11.00%
Calcium (Ca), Min.85%, Max.....	1.35%
Phosphorus (P), Minimum.....	0.45%
Potassium, Minimum.....	1.00%
Magnesium, Minimum.....	0.25%
Salt (NaCl), Min. 0.35%, Max.....	0.85%
Magnesium (Mg), Minimum.....	0.20%
Selenium (Se), Minimum.....	0.60 ppm
Vitamin A, Minimum.....	8,500 U/lb
Vitamin D, Minimum.....	2,700 IU/lb
Vitamin E, Minimum.....	71 IU/lb
Direct-Fed Microorganisms, Min....	334 million CFU/lb

INGREDIENTS

Cracked Corn, Dehulled Soybean Meal, Oats, Cottonseed Hulls, Distillers Dried Grains, Soybean Hulls, Beet Pulp, Wheat Middlings, Salt, Vitamin E Supplement, Ferrous Sulfate, Vitamin D Supplement, Manganese Sulfate, Zinc Sulfate, Manganese Amino Acid Complex, Zinc Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Selenium Yeast, Choline Chloride, Vitamin K Supplement, Copper Sulfate, Sodium Selenite, Vitamin A Supplement, Vitamin B12 supplement, Niacin Supplement, Iodine, d-Calcium Pantothenate, Biotin, Folic Acid, Riboflavin Supplement, Thiamine Mononitrate, Pyridoxine Hydrochloride, and Cobalt Sulfate.

FEEDING DIRECTIONS

Feed Rowe 18% Calf Start Sweet R30 to provide 0.14 -1.0 mg Monensin per pound of body weight per day to provide no more than 200 mg per head per day.

CAUTION: Do not use in feeds containing bentonite.

Warning: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin could be fatal to cattle. Must be thoroughly mixed in feeds before use. Do not exceed levels of Monensin recommended in the feeding directions, as reduced average daily gains may result

SIDNEY'S 18% CALF START

R30 BD PELLET

Complete Pellet for calves from birth to 444 pounds



Bovamine® Defend™ is a unique living probiotic that stabilizes rumen and small intestine digestive functions while stimulating natural and healthy digestion by reinforcing and enhancing the animal's natural defenses against harmful bacteria.

MEDICATED

For the prevention and control of coccidiosis due to *Eimeria bovis* and *E. Zuernii* and improved feed efficiency.

ACTIVE DRUG INGREDIENT

Monensin..... 30 Gms/Ton

Contains diflubenzuron

An insect growth regulator, which prevents the development of house, stable, face and horn flies in the manure of treated cattle

Diflubenzuron..... 3.65 Gms/Ton

GUARANTEED ANALYSIS

Crude Protein, Minimum.....	18.00%
Crude Fat, Minimum.....	4.00%
Crude Fiber, Maximum.....	9.50%
Acid Detergent Fiber (ADF), Maximum.....	12.00%
Calcium (Ca), Minimum.....	2.15%
Calcium (Ca), Maximum.....	2.65%
Phosphorus (P), Minimum.....	0.60%
Salt (NaCl), Minimum.....	0.50%
Salt (NaCl), Maximum.....	1.00%
Potassium (K), Minimum.....	0.95%
Selenium (Se), Minimum.....	.60 ppm
Vitamin A Minimum.....	8,600 IU/lb
Vitamin D Minimum.....	2,800 IU/lb
Vitamin E Minimum.....	70 IU/lb
Direct-Fed Microorganisms, Min.....	334 million CFU/lb

INGREDIENTS

Wheat Middlings, Corn Distillers Dried Grains, Ground Corn, Dehulled Soybean Meal, Calcium Carbonate, Cane Molasses, Salt, Montmorillonite Clay, Calcium Carbonate, Lignin Sulfonate, L-Lysine, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Vitamin E Supplement, Iron Sulfate, Vitamin D3 Supplement, Manganese Sulfate, Zinc Sulfate, Organic Selenium Yeast, Choline Chloride, Vitamin K Supplement, Corn Oil, Copper Sulfate, Sodium Selenite, Vitamin A Acetate in Gelatin, Vitamin B12 Supplement, Niacin, Iodine, Calcium Pantothenate, Biotin, Folic Acid, Riboflavin, Thiamine, Pyridoxine Hydrochloride, Cobalt Sulfate, Lactobacillus acidophilus and Propionibacterium freudenreichii.

FEEDING DIRECTIONS

Feed at the rate of 2.5 to 3% of bodyweight to provide .375 to .45 mg Monensin per pound of bodyweight not to exceed 200 mg (13.333 lb of feed) per head per day. and .10 mg/kg Diflubenzuron per day. Provide adequate amounts of roughage (pasture or hay). Consult Rowe Nutrition LLC for questions or more specific feeding instructions.

CAUTION: Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feed is safe for use in cattle only. Do not feed to lactating dairy cows. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as a reduced average daily gains may result. Feeding undiluted or mixing errors resulting in high concentrations of Monensin could be fatal to cattle. Must be thoroughly mixed in feeds before use. Do not feed undiluted. **WARNING:** A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

WARNING: Keep out of reach of children. Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling. For personal protective equipment (PPE), mixers and handlers must wear: long-sleeved shirt and pants, shoes and socks and chemical resistant gloves. User safety requirements are to follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. User safety recommendations are that users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and out on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. This pesticide is toxic to fish. Do not contaminate water when disposing of equipment wash



10107 US Route 127 North
West Manchester, Ohio
Office: 937-678-9015
Email: info@rowenutrition.com
Web: www.rowenutrition.com