



ROWE
Cattle

18% CALF STARTER R30 SWEET

Starter Feed for Calves up to 200 pounds

MEDICATED

For the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT

Monesin..... 30 gms/ton
Diflubenzuron..... 3.65 gms/ton

CONTAINS DIFLUBENZURON: An insect growth regulator, which prevents the development of house, stable, face, and horn flies in the manure of treated cattle.

GUARANTEED ANALYSIS

Crude Protein, Minimum.....	18.00%
Crude Fat, Minimum.....	2.50%
Crude Fiber, Maximum.....	11.00%
Calcium (Ca), Min.85%, Max.....	1.35%
Phosphorus (P), Minimum.....	0.45%
Potassium, Minimum.....	1.00%
Magnesium, Minimum.....	0.25%
Salt (NaCl), Min. 0.35%, Max.....	0.85%
Magnesium (Mg), Minimum.....	0.20%
Selenium (Se), Minimum.....	0.60 ppm
Vitamin A, Minimum.....	8,500 U/lb
Vitamin D, Minimum.....	2,700 IU/lb
Vitamin E, Minimum.....	71 IU/lb
Direct-Fed Microorganisms, Min...	334 million CFU/lb

INGREDIENTS

Cracked Corn, Dehulled Soybean Meal, Oats, Cottonseed Hulls, Distillers Dried Grains, Soybean Hulls, Beet Pulp, Wheat Middlings, Salt, Vitamin E Supplement, Ferrous Sulfate, Vitamin D Supplement, Manganese Sulfate, Zinc Sulfate, Manganese Amino Acid Complex, Zinc Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Selenium Yeast, Choline Chloride, Vitamin K Supplement, Copper Sulfate, Sodium Selenite, Vitamin A Supplement, Vitamin B12 supplement, Niacin Supplement, Iodine, d-Calcium Pantothenate, Biotin, Folic Acid, Riboflavin Supplement, Thiamine Mononitrate, Pyridoxine Hydrochloride, and Cobalt Sulfate.

FEEDING DIRECTIONS

Feed Rowe 18% Calf Start Sweet R30 to provide 0.14 -1.0 mg Monensin per pound of body weight per day to provide no more than 200 mg per head per day.

CAUTION: Do not use in feeds containing bentonite.

CAUTION: Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feed is safe for use in cattle only. Do not feed to lactating dairy cows. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as a reduced average daily gain may result. Feeding undiluted or mixing errors resulting in high concentrations of Monensin could be fatal to cattle. Must be thoroughly mixed in feeds before use. Do not feed undiluted. **WARNING:** A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.



10107 US Route 127 North West
Manchester, Ohio
Office: 937-678-9015
Email: info@rowenutrition.com
www.rowenutrition.com